

Reversible electrochemical capture and release of CO₂ using anthraquinone

Dominik Wielend(dominik.wielend@jku.at)¹, Dogukan Hazar Apaydin², Dong Ryeol Whang¹

Niyazi Serdar Sariciftci¹

¹Linz Institute for Organic Solar Cells (LIOS), Institute of Physical Chemistry, Johannes Kepler University Linz, Altenbergerstrasse 69, 4040 Linz, Austria., ²Linz Institute for Organic Solar Cells (LIOS), Institute of Physical Chemistry, Johannes Kepler University Linz, Altenbergerstrasse 69, 4040 Linz, Austria.; Current address: Institute of Science and Technology Austria (IST Austria), Am Campus 1, 3400 Klosterneuburg, Austria.

Abstract

The general strategy of Carbon Capture and Utilization (CCU) is the usage of carbon dioxide (CO₂) as feedstock material for sustainable and fossil-fuel free chemicals. An important requirement prior catalysing the conversion of CO₂ is a controlled capturing and release process of CO₂ [1]. As the industrial absorption processes require either high heat or vacuum input, developing more benign alternatives is still a challenging subject [2].

Recent examples of technologies for electrochemical capture of CO₂ are using organic molecules like sulfides [3], bipyridines [4,5] or carbonyl compounds [6], mostly homogeneously dissolved in organic solutions. In order to circumvent the use of large quantities of organic solvents, our group recently reported electrochemical adsorption and desorption of CO₂ in an ambient condition by aromatic carbonyl pigments, like quinacridone [7] or a naphthalene bisimide derivatative [8].

As a continuing series of the previous works, herein we demonstrate that anthraquinone thin-film electrodes can electrochemically capture and release CO₂ with an uptake capacity of 5.9 mmol g⁻¹, which is comparable with that of an industrial amine process (8 mmol g⁻¹) [8,9]. In contrast to the previous studies, anthraquinone is industrially used and therefore cheap and moreover we could demonstrate this process to fully operate in aqueous solution. Furthermore, the proposed chemical formation of anthraquinone carbonate-like structure was supported by in-depth studies using in-situ spectroelectrochemistry with UV-vis and ATR-FTIR techniques [10].

Keywords: organic pigments, electrochemistry, carbon dioxide capture, spectroelectrochemistry

References

- [1] M. Aresta, Carbon Dioxide as Chemical Feedstock, WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim, 2010.
- [2] S. Topham, A. Bazzanella, S. Schiebahn, S. Luhr, L. Zhao, A. Otto and D. Stolten, Ullmann's Encycl. Ind. Chem., 2014, 1–43.
- [3] P. Singh, J. H. Rheinhardt, J. Z. Olson, P. Tarakeshwar, V. Mujica and D. A. Buttry, J. Am. Chem. Soc., 2017, 139, 1033–1036.
- [4] H. Ishida, T. Ohba, T. Yamaguchi and K. Ohkubo, Chem. Lett., 1994, 23, 905–908.
- [5] R. Ranjan, J. Olson, P. Singh, E. D. Lorange, D. A. Buttry and I. R. Gould, J. Phys. Chem. Lett., 2015, 6, 4943–4946.
- [6] B. Gurkan, F. Simeon and T. A. Hatton, ACS Sustain. Chem. Eng., 2015, 3, 1394–1405.
- [7] D. H. Apaydin, E. D. Głowacki, E. Portenkirchner and N. S. Sariciftci, Angew. Chemie - Int. Ed., 2014, 53, 6819–6822.
- [8] D. H. Apaydin, M. Gora, E. Portenkirchner, K. T. Oppelt, H. Neugebauer, M. Jakesova, E. D. Głowacki, J. Kunze-Liebhäuser, M. Zagorska, J. Mieczkowski and N. S. Sariciftci, ACS Appl. Mater. Interfaces, 2017, 9, 12919–12923.
- [9] T. Supap, R. Idem, P. Tontiwachwuthikul and C. Saiwan, Int. J. Greenh. Gas Control, 2009, 3, 133–142.
- [10] D. Wielend, D. H. Apaydin and N. S. Sariciftci, J. Mater. Chem. A, 2018, 6, 15095–15101.

Biography

Dominik Wielend graduated from the bachelor as well as the master studies in technical chemistry from the

Johannes Kepler University Linz in 2015 and 2017, respectively. In 2016/2017 Dominik Wielend did an Erasmus internship in the Christian Doppler Laboratory for Sustainable Syngas Chemistry / Erwin Reisner Lab at the University of Cambridge (United Kingdom).

The current research topic for the PhD studies involve the investigation of organic electro-catalysts for oxygen reduction.